

University of Groningen

Enantiocomplementary Epoxidation Reactions Catalyzed by an Engineered Cofactor-Independent Non-Natural Peroxygenase

Xu, Guangcai; Crotti, Michele; Thangavelu, Saravan; Kataja, Kim; Poelarends, Gerrit J.

Published in:
Angewandte Chemie International Edition

DOI:
[10.1002/anie.202001373](https://doi.org/10.1002/anie.202001373)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2020

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Xu, G., Crotti, M., Thangavelu, S., Kataja, K., & Poelarends, G. J. (2020). Enantiocomplementary Epoxidation Reactions Catalyzed by an Engineered Cofactor-Independent Non-Natural Peroxygenase. *Angewandte Chemie International Edition*, 59(26), 10374-10378. <https://doi.org/10.1002/anie.202001373>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Biocatalysis

How to cite: *Angew. Chem. Int. Ed.* **2020**, 59, 10374–10378

International Edition: doi.org/10.1002/anie.202001373

German Edition: doi.org/10.1002/ange.202001373

Enantiocomplementary Epoxidation Reactions Catalyzed by an Engineered Cofactor-Independent Non-natural Peroxygenase

Guangcai Xu, Michele Crotti, Thangavelu Saravanan, Kim M. Kataja, and Gerrit J. Poelarends*

Abstract: Peroxygenases are heme-dependent enzymes that use peroxide-borne oxygen to catalyze a wide range of oxyfunctionalization reactions. Herein, we report the engineering of an unusual cofactor-independent peroxxygenase based on a promiscuous tautomerase that accepts different hydroperoxides (*t*-BuOOH and H₂O₂) to accomplish enantiocomplementary epoxidations of various α,β -unsaturated aldehydes (citral and substituted cinnamaldehydes), providing access to both enantiomers of the corresponding α,β -epoxy-aldehydes. High conversions (up to 98 %), high enantioselectivity (up to 98 % ee), and good product yields (50–80 %) were achieved. The reactions likely proceed via a reactive enzyme-bound iminium ion intermediate, allowing tweaking of the enzyme's activity and selectivity by protein engineering. Our results underscore the potential of catalytic promiscuity for the engineering of new cofactor-independent oxidative enzymes.

Peroxygenases are oxidative enzymes catalyzing the insertion of an oxygen atom into a variety of substrates using peroxide-borne oxygen instead of dioxygen.^[1] Earlier studies have mainly focused on the peroxxygenase activity of chloroperoxidases that normally catalyze peroxide-driven halogenation of organic compounds^[2] and engineered cytochrome P450 monooxygenases that are able to promote peroxide-mediated hydroxylation reactions.^[3] However, since the discovery of the unspecific peroxxygenase *AaeUPO* in 2004,^[4] which uses hydrogen peroxide to catalyze diverse oxyfunctionalization reactions including oxidations, sulfoxidations, epoxidations and hydroxylations,^[5] there has been an increasing interest to discover and develop novel peroxxygenases. Note that peroxxygenases are cofactor-dependent enzymes, typically relying on heme to react with hydrogen

peroxide, forming an oxoferryl-heme as the actual oxygenating species to promote oxyfunctionalization reactions.^[1,5a,6]

In this study, we report the engineering of an unusual cofactor-independent peroxxygenase based on the promiscuous peroxxygenase activity of the enzyme 4-oxalocrotonate tautomerase (4-OT). This engineered non-natural peroxxygenase accepts different hydroperoxides (*t*-BuOOH and H₂O₂) to achieve enantiocomplementary epoxidations of various α,β -unsaturated aldehydes, providing access to both enantiomers of the resultant α,β -epoxy-aldehydes with high enantiomeric excess (up to 98 %). Our results showcase the potential of catalytic promiscuity for the creation of new cofactor-independent oxidative enzymes.

We have recently demonstrated that 4-OT can catalyze the Michael addition of nitromethane to cinnamaldehyde via an enzyme-bound iminium ion intermediate, which is formed between cinnamaldehyde and the catalytic active site Pro-1 residue.^[7] In this study, we examined if this catalytic mechanism of 4-OT could be exploited for other useful synthetic applications, and envisioned that substituting nitromethane with *t*-BuOOH could trigger a C–O bond formation between the reactive enzyme-bound iminium ion species and *t*-BuOOH (Supporting Information, Scheme S1). The resulting enamine intermediate may undergo ring closure to construct the final epoxide moiety. To our delight, when wild-type 4-OT (0.33 mg mL^{−1}) was incubated at room temperature with 200 mM *t*-BuOOH (**1**, Figure 1) and 1 mM cinnamaldehyde (**2a**) in 20 mM sodium phosphate buffer (pH 7.3) containing 5 % (v/v) EtOH, 38 % of substrate **2a** was consumed within 48 h and the corresponding α,β -epoxy-aldehyde **3a** was identified by GC-MS analysis. Chiral HPLC analysis of the alcohol derivative of **3a** revealed the product to be (2*R*,3*S*)-**3a** with an e.r. of 62:38 and a d.r. of 92:8 (*syn/anti*). The *S* configuration of the C3 carbon of **3a** indicates that *t*-BuOOH selectively attacks cinnamaldehyde from the *re*-face, similar to what was previously observed for the nitromethane addition reaction.^[7] These results demonstrated that 4-OT has low-level promiscuous peroxxygenase activity and may possibly serve as a template to construct a cofactor-independent peroxxygenase.

Motivated by these initial findings, a mutability-landscape-guided protein engineering approach^[8] was applied to enhance the promiscuous peroxxygenase activity of wild-type 4-OT. For this, a previously constructed collection of 4-OT genes encoding nearly all possible single-mutant variants of 4-OT^[9] was screened to identify hotspot positions in the protein where mutations give improved peroxxygenase activity. Improved variants were identified by monitoring the depletion of substrate **2a** in a spectrophotometric kinetic assay in multiwell plates (see Supporting Information). Interestingly,

[*] G. Xu, Dr. M. Crotti, Dr. T. Saravanan, K. M. Kataja, Prof. Dr. G. J. Poelarends
Department of Chemical and Pharmaceutical Biology, Groningen Research Institute of Pharmacy, University of Groningen
Antonius Deusinglaan 1, 9713 AV Groningen (The Netherlands)
E-mail: g.j.poelarends@rug.nl

Dr. T. Saravanan

Present address: School of Chemistry, University of Hyderabad
P.O. Central University, Gachibowli, Hyderabad 500046 (India)

Supporting information and the ORCID identification number(s) for the author(s) of this article can be found under:
<https://doi.org/10.1002/anie.202001373>.

© 2020 The Authors. Published by Wiley-VCH Verlag GmbH & Co. KGaA. This is an open access article under the terms of the Creative Commons Attribution Non-Commercial NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial, and no modifications or adaptations are made.




Figure 1. Peroxygenase activity of 4-OT variants. A) Reaction scheme of 4-OT-catalyzed epoxidation of **2a** using *t*-BuOOH as oxidant. B) Comparison of the peroxxygenase activity of wild-type 4-OT (WT) and engineered 4-OT variants. C) Reaction progress curves using different 4-OT variants. The initial rate of the 4-OT(YIA)-catalyzed reaction corresponds to 16 $\mu\text{M min}^{-1}$; specific activity is 48 mU per mg of protein.

six mutations at position F50 significantly improved (> 5-fold) the peroxxygenase activity, with F50A (10-fold enhanced activity) and F50V (25-fold enhanced activity) being the best mutant enzymes (Figure 1). Several mutations at positions Q4, I41, M45, and H49 also resulted in enzyme variants with 2- to 5-fold enhanced activity compared to wild-type 4-OT, including mutants Q4Y, I41Y, M45I, and H49T. The most beneficial mutations, F50V and F50A, were subsequently combined with mutations at other hotspot positions by using site-saturation mutagenesis (Table S2). Activity screening of the seven mutant libraries resulted in the identification of three double mutants (Q4Y/F50V, Q4F/F50A and M45I/F50A) with around 1.6- to 3.5-fold enhanced activity compared to the parental enzymes. Based on these results, the mutant libraries Q4X/M45I/F50A, Q4Y/M45X/F50A, Q4F/M45X/F50A, and Q4Y/M45I/F50X were constructed (Table S2). Screening of these four libraries led to the discovery of the triple mutant Q4Y/M45I/F50A (YIA) that showed a remarkable 60-fold enhancement in activity for the epoxidation of **2a** compared to the wild-type enzyme (Figure 1). Notably, this mutant enzyme also has enhanced enantioselectivity, allowing the production of (2*R*,3*S*)-**3a** with

high enantiopurity (e.r. = 98:2). Replacement of Pro-1 with an alanine in 4-OT(YIA) led to a 210-fold decrease in *t*-BuOOH-dependent epoxidation of **2a** (Figure S31), providing support for a reaction mechanism that proceeds via iminium ion formation between the active site Pro-1 residue and the α,β -unsaturated aldehyde substrate (Scheme S1).

With the improved 4-OT mutant YIA in hand, the substrate scope of this enzyme was explored by testing a set of α,β -unsaturated aldehydes using *t*-BuOOH as oxidant. The results showed that 4-OT(YIA) has a broad substrate scope, accepting *ortho*-, *meta*-, and *para*-substituted cinnamaldehydes (**2a–i**, Figure 2), and catalyzes their epoxidation with good conversions (94–97%) to yield the corresponding products (**3a–i**) with excellent enantiopurity (e.r. up to 99:1). The d.r. values (up to 92:8) obtained indicate that the enzyme has good stereocontrol over both carbons of the double bond of **2**. The enzyme also accepts the aliphatic α,β -unsaturated aldehyde citral (**2j**) with good enantioselectivity, moderate diastereoselectivity, and excellent regioselectivity (Figure 2). All epoxide products have the *syn* configuration with 2*R*,3*S* being the major enantiomer, indicating that *t*-BuOOH exclusively attacked from the *re*-face of the enzyme-bound iminium ion intermediate (Figure 3). Note that while the enzymatic α,β -epoxy-aldehyde products **3** have a 2*R*,3*S* configuration, the corresponding α,β -epoxy-alcohols have a 2*S*,3*S* configuration. This deviant configuration of the α,β -epoxy-alcohols is due to different prioritization of the substituents at the C2 chiral center relative to their α,β -epoxy-aldehyde precursors.

Given that *t*-BuOOH is a relatively large hydroperoxide, we investigated if the use of a small hydroperoxide such as H_2O_2 would change the preference of attack to the *si*-face (Figure 3). Again, compound **2a** was used as a substrate for initial testing with H_2O_2 as oxidant and mutant 4-OT(YIA) as catalyst. Notably, the epoxidations with H_2O_2 showed relatively high non-enzymatic (background) reaction rates when applying the same conditions as used with *t*-BuOOH. To suppress the background reaction, without diminishing the enzymatic activity, the pH of the reaction was lowered from 7.3 to 6.3 (Figure S1). Gratifyingly, the results showed that the use of H_2O_2 not only inverted the stereochemistry of the reaction, but product **3a** was also obtained with good e.r. (92:8) and d.r. (84:16) values (Figure 2 and Figure 3). Thus, an enantiocomplementary epoxidation of **2a** was achieved by replacing *t*-BuOOH by H_2O_2 as the oxidant. Next, we evaluated the same set of α,β -unsaturated aldehydes **2b–2j** for 4-OT(YIA)-catalyzed epoxidation reactions using H_2O_2 (Figure 2). Importantly, the stereo inversions occurred with all aromatic substrates (**2a–2i**), and high conversions and good enantio- and diastereoselectivity were observed. Notably, the epoxidation reaction with the aliphatic substrate citral (**2j**) also showed an inversion of diastereoselectivity when using H_2O_2 instead of *t*-BuOOH, resulting in the *anti*-product as the major isomer; the *syn*-product also showed an inverted enantiomer ratio (Figure S30).

Having achieved enantiocomplementary epoxidation reactions, we next investigated semi-preparative scale chemoenzymatic synthesis of α,β -epoxy-alcohols **4a–j** using 4-OT(YIA) as catalyst. With *t*-BuOOH as oxidant, the reactions




Figure 2. Substrate scope of the epoxidation reactions catalyzed by 4-OT(YIA) (Q4Y/M45I/F50A) using *t*-BuOOH or H₂O₂ as the oxidant. Reaction conditions: **2** (1 mM), *t*-BuOOH (100 mM) or H₂O₂ (25 mM), 4-OT(YIA) (0.1 mg mL⁻¹), NaPi buffer (20 mM), 5% (v/v) EtOH (MeCN for **2f**, **2g** and **2h**). Conversions were calculated based on the depletion of the absorbance corresponding to substrate **2**. Product identifications were performed by GC-MS analysis. The enzymatic α,β -epoxy-aldehyde products **3** were reduced to the corresponding α,β -epoxy-alcohols with NaBH₄ for chiral HPLC analysis. The e.r., d.r. and absolute stereochemistry of products were determined by chiral HPLC using authentic standards. The diastereomer ratio (d.r.) is defined as the *syn/anti* ratio. **2j**: *E/Z* = 3:2; n.d.: not determined.

were performed with higher substrate concentrations (50 mM **2a**, 20 mM **2b–g** and **2j**, 5 mM **2i**), maintaining high conversions and stereoselectivity with good isolated product yields (Table 1). Only a minor decrease in enantio- and

diastereoselectivity was observed, most likely due to the non-enzymatic background reaction, since somewhat longer reaction times were needed. The semi-preparative scale synthesis using H₂O₂ as oxidant was also attempted using

group is focused on further improving the peroxygenase activity of this promising cofactor-independent enzyme, making use of a recently developed selective colorimetric “turn-on” probe for efficient engineering of iminium biocatalysis.^[13] The non-natural peroxygenase activity of 4-OT, together with its previously reported promiscuous C–C bond-forming Michaelase^[7,9,14] and aldolase^[15] activities, emphasize the chemical versatility of the 4-OT protein scaffold.

Acknowledgements

We acknowledge financial support from the Netherlands Organization of Scientific Research (VICI grant 724.016.002 and ECHO grant 713.015.003).

Conflict of interest

The authors declare no conflict of interest.

Keywords: enzyme engineering · epoxidation · oxidative enzymes · peroxide · peroxygenase

- [1] Y. Wang, D. Lan, R. Durrani, F. Hollmann, *Curr. Opin. Chem. Biol.* **2017**, *37*, 1–9.
- [2] a) D. R. Morris, L. P. Hager, *J. Biol. Chem.* **1966**, *241*, 1763–1768; b) A. Zaks, D. R. Dodds, *J. Am. Chem. Soc.* **1995**, *117*, 10419–10424; c) J. Littlechild, *Curr. Opin. Chem. Biol.* **1999**, *3*, 28–34.
- [3] a) H. Joo, Z. Lin, F. H. Arnold, *Nature* **1999**, *399*, 670–673; b) P. C. Cirino, F. H. Arnold, *Angew. Chem. Int. Ed.* **2003**, *42*, 3299–3301; *Angew. Chem.* **2003**, *115*, 3421–3423; c) J. Chen, F. Kong, N. Ma, P. Zhao, C. Liu, X. Wang, Z. Cong, *ACS Catal.* **2019**, *9*, 7350–7355.
- [4] R. Ullrich, J. Nueske, K. Scheibner, J. Spantzel, M. Hofrichter, *Appl. Environ. Microbiol.* **2004**, *70*, 4575–4581.
- [5] a) M. Hofrichter, R. Ullrich, *Curr. Opin. Chem. Biol.* **2014**, *19*, 116–125; b) I. Bassanini, E. E. Ferrandi, M. Vanoni, G. Ottolina, S. Riva, M. Crotti, E. Brenna, D. Monti, *Eur. J. Org. Chem.* **2017**, 7186–7189.
- [6] a) C. K. Prier, F. H. Arnold, *J. Am. Chem. Soc.* **2015**, *137*, 13992–14006; b) F. H. Arnold, *Angew. Chem. Int. Ed.* **2018**, *57*, 4143–4148; *Angew. Chem.* **2018**, *130*, 4212–4218.
- [7] C. Guo, M. Saifuddin, T. Saravanan, G. J. Poelarends, *ACS Catal.* **2019**, *9*, 4369–4373.
- [8] J. Y. van der Meer, L. Biewenga, G. J. Poelarends, *ChemBioChem* **2016**, *17*, 1792–1799.
- [9] J. Y. Van Der Meer, H. Poddar, B. J. Baas, Y. Miao, M. Rahimi, A. Kunzendorf, R. Van Merkerk, P. G. Tepper, E. M. Geertsema, A. M. W. H. Thunnissen, et al., *Nat. Commun.* **2016**, *7*, 1–16.
- [10] a) P. Zhou, X. Wang, B. Yang, F. Hollmann, Y. Wang, *RSC Adv.* **2017**, *7*, 12518–12523; b) S. Ranganathan, S. Zeitlhofer, V. Sieber, *Green Chem.* **2017**, *19*, 2576–2586.
- [11] M. Svedendahl, P. Carlqvist, C. Branneby, O. Allnér, A. Frise, K. Hult, P. Berglund, T. Brinck, *ChemBioChem* **2008**, *9*, 2443–2451.
- [12] a) S. Juliá, J. Masana, J. C. Vega, *Angew. Chem. Int. Ed. Engl.* **1980**, *19*, 929–931; *Angew. Chem.* **1980**, *92*, 968–969; b) S. Colonna, A. Manfredi, *Tetrahedron Lett.* **1986**, *27*, 387–390; c) S. Colonna, A. Manfredi, R. Annunziata, M. Spadoni, *Tetrahedron* **1987**, *43*, 2157–2164; d) S. Baars, K.-H. Drauz, H.-P. Krimmer, S. M. Roberts, J. Sander, J. Skidmore, G. Zanardi, *Org. Process Res. Dev.* **2003**, *7*, 509–513; e) H. Sundén, I. Ibrahim, A. Córdova, *Tetrahedron Lett.* **2006**, *47*, 99–103.
- [13] L. Biewenga, M. Crotti, M. Saifuddin, G. J. Poelarends, *ACS Omega* **2020**, *5*, 2397–2405.
- [14] a) E. Zandvoort, E. M. Geertsema, B. J. Baas, W. J. Quax, G. J. Poelarends, *Angew. Chem. Int. Ed.* **2012**, *51*, 1240–1243; *Angew. Chem.* **2012**, *124*, 1266–1269; b) Y. Miao, E. M. Geertsema, P. G. Tepper, E. Zandvoort, G. J. Poelarends, *ChemBioChem* **2013**, *14*, 191–194; c) E. M. Geertsema, Y. Miao, P. G. Tepper, P. Dehaan, E. Zandvoort, G. J. Poelarends, *Chem. Eur. J.* **2013**, *19*, 14407–14410; d) Y. Miao, P. G. Tepper, E. M. Geertsema, G. J. Poelarends, *Eur. J. Org. Chem.* **2016**, 5350–5354; e) L. Biewenga, T. Saravanan, A. Kunzendorf, J. Van Der Meer, T. Pijning, P. G. Tepper, R. Van Merkerk, S. J. Charnock, A. W. H. Thunnissen, G. J. Poelarends, *ACS Catal.* **2019**, *9*, 1503–1513.
- [15] a) E. Zandvoort, B. J. Baas, W. J. Quax, G. J. Poelarends, *ChemBioChem* **2011**, *12*, 602–609; b) M. Rahimi, J. Y. van der Meer, E. M. Geertsema, H. Poddar, B. J. Baas, G. J. Poelarends, *ChemBioChem* **2016**, *17*, 1225–1228; c) M. Rahimi, E. M. Geertsema, Y. Miao, J.-Y. van der Meer, T. van den Bosch, P. de Haan, E. Zandvoort, G. J. Poelarends, *Org. Biomol. Chem.* **2017**, *15*, 2809–2816.

Manuscript received: January 27, 2020

Revised manuscript received: March 6, 2020

Accepted manuscript online: March 11, 2020

Version of record online: April 14, 2020